Fluid Phase Equilibria, 50 (1989) 349–350
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 50 (1989)

Volume 50, Nos. 1–2

On the application of cubic equations of state: analytical expression for α/T_r improved liquid density calculations
H. Sugie, Y. Iwahori (Nagoya, Japan) and B.C.-Y. Lu (Ottawa, Ont., Canada) 1– 20

Extension of the AEOS model to systems containing any number of associating and inert components
A. Anderko (Warsaw, Poland) 21– 52

The generalized van der Waals partition function. V. Mixture of square-well fluids of different sizes and energies
K.-H. Lee (Pohang, Korea), L.R. Dodd and S.I. Sandler (Newark, DE, U.S.A.) 53– 77

Phase equilibria in mixtures of glycerides and carbon dioxide and application of continuous thermodynamics to mixtures of rapeseed oil and carbon dioxide
T. Klein and S. Schulz (Dortmund, F.R.G.) 79–100

Liquid–liquid equilibria for binary and ternary systems including the critical region. Transformation to non-classical coordinates
J.J. De Pablo and J.M. Prausnitz (Berkeley, CA, U.S.A.). 101–131

Salt effect in phase equilibria: effect of dissolved inorganic salts on the liquid–liquid equilibria of benzene–2-propanol–water system and the vapor–liquid equilibria of its constituent binaries
M. Rajendran, S. Renganarayanan and D. Srinivasan (Madras, India) 133–164

Liquid–liquid–vapor phase equilibria behavior of certain binary nitrous oxide + n-alkylbenzene mixtures
D.H. Lam, A. Jangkamolkulchai and K.D. Luks (Tulsa, OK, U.S.A.) 165–174

Multiphase equilibria behavior of nitrous oxide + n-paraffin mixtures
A. Jangkamolkulchai, D.H. Lam and K.D. Luks (Tulsa, OK, U.S.A.) 175–187

Reexamination of the multiphase equilibria of the system carbon dioxide + n-butylbenzene + n-eicosane
M.M. Miller, A. Jangkamolkulchai and K.D. Luks (Tulsa, OK, U.S.A.) 189–199

Vapor–liquid phase equilibrium measurements for the n-hexane–1-butanol system at 323.15, 338.15 and 348.15 K
M.R.W. Maciel and A.Z. Francesconi (Campinas, Brazil) 201–208

Infinite-dilution activity coefficients of water in TEG, PEG, glycerol and their mixtures in the temperature range 50 to 140 °C
B. Bestani and K.S. Shing (Los Angeles, CA, U.S.A.) 209–221

Solubility of He, Ne, Ar, Kr, Xe, H_2, D_2, N_2, O_2, CH_4, C_2H_4, C_2H_6, CF_4, SF_6 and CO_2 in cyclopentanone from 273.15 K to 303.15 K and gas partial pressure of 101.33 kPa
M.A. Gallardo, M.C. López, J.S. Urieta and C. Gutiérrez Losa (Zaragoza, Spain) 223–233

Volume 50, No. 3

Excess volumes of binary systems formed by a pyridine base and an n-alkane in terms of an association model
A.J. Treszczanowicz and D. Patterson (Montreal, P.Q., Canada), G.C. Benson (Ottawa, Ont., Canada) and T. Kasprzycka-Guttman (Warsaw, Poland) 235–247

Application of a maximum likelihood method using implicit constraints to determine equation of state parameters from binary phase behavior data
V.G. Niesen and V.F. Yesavage (Golden, CO, U.S.A.) 249–266

Excess properties of mixtures of polar components and hydrocarbons of varying local polarisability
N.V. Nhu and F. Kohler (Bochum, F.R.G.) 267–296

Excess volumes for binary liquid mixtures of methylethylketone with methylene chloride, 1,2-dichloroethane, trichloroethylene, tetrachloroethylene and cyclohexane at various temperatures
J. Nath and R. Saini (Gorakhpur, India) 297–303

Phase equilibria in binary mixtures of ethane and hexadecane
R. de Goede, C.J. Peters, H.J. van der Kooi and R.N. Lichtenthaler (Delft, The Netherlands) 305–314

Vapor–liquid equilibria for binary mixtures of carbon dioxide and fatty acid ethyl esters
R. Bharath, H. Inomata, K. Arai, K. Shoji and Y. Noguchi (Sendai, Japan) .. 315–327

Vapour–liquid equilibria of binary systems of C_1–C_4 *n*-alcohols with 1,2-dichloroethane at 323.15 K
S.K. Chaudhari and S.S. Katti (Poona, India) 329–338

Liquid–liquid equilibrium in the toluene–methyl ethyl ketone–water system
R.C. Ghildiyal, S.M. Nanoti, N.N. Kulsrestha, B.S. Rawat, Y. Kumar and R. Krishna (Dehra Dun, India) 339–346

Author Index 347

Contents of Volume 50 (1989) 349–350